

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:SSPTASXS1656

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * * * * * Welcome to STN International * * * * *

NEWS 1 Web Page for STN Seminar Schedule - N. America
NEWS 2 AUG 10 Time limit for inactive STN sessions doubles to 40 minutes
NEWS 3 AUG 18 COMPENDEX indexing changed for the Corporate Source (CS) field
NEWS 4 AUG 24 ENCOMPLIT/ENCOMPLIT2 reloaded and enhanced
NEWS 5 AUG 24 CA/CAplus enhanced with legal status information for U.S. patents
NEWS 6 SEP 09 50 Millionth Unique Chemical Substance Recorded in CAS REGISTRY
NEWS 7 SEP 11 WPIDS, WPINDEX, and WPIX now include Japanese FTERM thesaurus
NEWS 8 OCT 21 Derwent World Patents Index Coverage of Indian and Taiwanese Content Expanded
NEWS 9 OCT 21 Derwent World Patents Index enhanced with human translated claims for Chinese Applications and Utility Models
NEWS 10 NOV 23 Addition of SCAN format to selected STN databases
NEWS 11 NOV 23 Annual Reload of IFI Databases
NEWS 12 DEC 01 FRFULL Content and Search Enhancements
NEWS 13 DEC 01 DGENE, USGENE, and PCTGEN: new percent identity feature for sorting BLAST answer sets
NEWS 14 DEC 02 Derwent World Patent Index: Japanese FI-TERM thesaurus added
NEWS 15 DEC 02 PCTGEN enhanced with patent family and legal status display data from INPADOCDB
NEWS 16 DEC 02 USGENE: Enhanced coverage of bibliographic and sequence information
NEWS 17 DEC 21 New Indicator Identifies Multiple Basic Patent Records Containing Equivalent Chemical Indexing in CA/CAplus

NEWS EXPRESS MAY 26 09 CURRENT WINDOWS VERSION IS V8.4,
AND CURRENT DISCOVER FILE IS DATED 06 APRIL 2009.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN customer agreement. This agreement limits use to scientific research. Use for software development or design, implementation of commercial gateways, or use of CAS and STN data in the building of commercial

products is prohibited and may result in loss of user privileges and other penalties.

FILE 'HOME' ENTERED AT 23:42:18 ON 22 DEC 2009

FILE 'MEDLINE' ENTERED AT 23:42:31 ON 22 DEC 2009

FILE 'SCISEARCH' ENTERED AT 23:42:31 ON 22 DEC 2009
Copyright (c) 2009 The Thomson Corporation

FILE 'LIFESCI' ENTERED AT 23:42:31 ON 22 DEC 2009
COPYRIGHT (C) 2009 Cambridge Scientific Abstracts (CSA)

FILE 'BIOTECHDS' ACCESS NOT AUTHORIZED

FILE 'BIOSIS' ENTERED AT 23:42:31 ON 22 DEC 2009
Copyright (c) 2009 The Thomson Corporation

FILE 'EMBASE' ENTERED AT 23:42:31 ON 22 DEC 2009
Copyright (c) 2009 Elsevier B.V. All rights reserved.

FILE 'HCAPLUS' ENTERED AT 23:42:31 ON 22 DEC 2009
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'NTIS' COULD NOT BE ENTERED

FILE 'ESBIOBASE' COULD NOT BE ENTERED

FILE 'BIOTECHNO' COULD NOT BE ENTERED

FILE 'WPTDS' COULD NOT BE ENTERED

=> S Clupeine or protamine (p) (ompt protease or protease VII)
L1 685 CLUPEINE OR FROTAMINE (P) (OMPT PROTEASE OR PROTEASE VII)

=> S Clupeine (p) (ompt protease or protease VII)
L2 0 CLUPEINE (P) (OMPT PROTEASE OR PROTEASE VII)

=> S protamine (4a) (ompt protease or protease VII)
L3 1 PROTAMINE (4A) (OMPT PROTEASE OR PROTEASE VII)

=> d 13 bib ab

I-3 ANSWER 1 OF 1 HCAPLUS COPYRIGHT 2009 ACS on STN

AN 1998-524254 HCAPLUS

DN 129·214014

QBEF 129:43415a 43418a

TI Identification of OmpT as the protease that hydrolyzes the antimicrobial peptide protamine before it enters growing cells of *Escherichia coli*

All Peptides prolineamide before it effects growing cells of Escherichia coli
Stumper, Stefan; Schmidt, Roland; Stephens, Darren L.; Georgiou, George:

Bakker, Evert P.
CS Abteilung Mikrobiologie, Universitat Osnabruck, Osnabruck, D-49069,
Germany
SO Journal of Bacteriology (1998), 180(15), 4002-4006
CODEN: JOBAAY; ISSN: 0021-9193
PB American Society for Microbiology
DT Journal
LA English
AB The influence of extracytoplasmic proteases on the resistance of *Escherichia coli* to the antimicrobial peptide protamine was investigated by testing strains with deletions in the protease genes *degP*, *ptr*, and *ompT*. Only Δ *ompT* strains were hypersusceptible to protamine. This effect was abolished by plasmids carrying *ompT*. Both at low and at high Mg²⁺ concns., *ompT*⁺ strains cleared protamine from the medium within a few minutes. By contrast, at high Mg²⁺ concns., protamine remained present for at least 1 h in the medium of an *ompT* strain. These data indicate that OmpT is the protease that degrades protamine and that it exerts this function at the external face of the outer membrane.
OSC.G 86 THERE ARE 86 CAPLUS RECORDS THAT CITE THIS RECORD (86 CITINGS)
RE.CNT 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT